

**Table 1.** Summary of findings from interviews with post-overdose outreach staff and managers (n=21), organized by CFIR domains

---

<b>Stimulant Use Landscape and Patterns of Use (CFIR Outer Setting)</b>
Types of Stimulants that Overdose Survivors are Using
<ul style="list-style-type: none"><li>• Crack and cocaine common; methamphetamine use increasing in rural areas</li><li>• Stimulant use was rarely isolated; often combined with fentanyl/xylazine</li></ul>
Circumstances and Settings of Stimulant Use Among Survivors
<ul style="list-style-type: none"><li>• Social/recreational (“weekend warriors”), chemsex, and functional use (energy, staying awake, safety for unhoused)</li></ul>
<b>Identification of Stimulant-Involved Events (CFIR Inner Setting)</b>
First Responder Emergency Call Logs
<ul style="list-style-type: none"><li>• Heavy reliance on 911/police/EMS data for identification; quality inconsistent</li></ul>
Challenges Identifying Stimulant-Involved Events
<ul style="list-style-type: none"><li>• Stimulant-involved overdoses and related acute health events often missed: reliance on naloxone administration, misclassified as mental health, no 911 call</li></ul>
Expanding and Improving Identification of Stimulant-Involved Events
<ul style="list-style-type: none"><li>• Word-of-mouth and first responder training as opportunities</li></ul>
<b>Perceived Effectiveness and Capacity Building Needs (CFIR Inner Setting)</b>
Perceived Effectiveness of Post-Overdose Outreach for Stimulant-Involved Events
<ul style="list-style-type: none"><li>• Outreach perceived as less effective for stimulants than opioids</li></ul>
Capacity Building Needs for Post-Overdose Outreach Programs
<ul style="list-style-type: none"><li>• Need for stimulant-specific training on recognition and response</li></ul>
<b>Enhancing Post-Overdose Outreach for Stimulant-Involved Events (CFIR Process)</b>
Awareness and Knowledge Building
<ul style="list-style-type: none"><li>• Low awareness of risk and perceived susceptibility; stimulants deprioritized vs. opioids</li></ul>
Harm Reduction Supplies and Safer Use Supports
<ul style="list-style-type: none"><li>• Smoking/snorting kits, fentanyl test strips, and drug checking as engagement tools</li></ul>
Specialized Settings and Contexts
<ul style="list-style-type: none"><li>• Expand outreach to more non-traditional community settings; limited evening capacity</li></ul>
Referral and Navigation
<ul style="list-style-type: none"><li>• Few stimulant-specific services; major provider barriers</li></ul>

---